



Thomas, A., Bailey, M., Finn, A., Fleming, H., Lee, M., Mead, A., Morales-Aza, B., Vipond, I., & Eisler, M. (2018). *Attenuated vaccines as model respiratory viral infection systems: from human to bovine*. Paper presented at UK Veterinary Vaccinology Network Conference 2018, Stirling, Scotland, United Kingdom.

Peer reviewed version

[Link to publication record in Explore Bristol Research](#)
PDF-document

University of Bristol - Explore Bristol Research

General rights

This document is made available in accordance with publisher policies. Please cite only the published version using the reference above. Full terms of use are available:
<http://www.bristol.ac.uk/red/research-policy/pure/user-guides/ebr-terms/>

Attenuated vaccines as model respiratory viral infection systems: from human to bovine

A. Thomas^{1,2,4}, M. Bailey¹, A. Finn⁴, H.R. Fleming², M.R.F. Lee^{1,2}, A. Mead³,
B. Morales-Aza⁴, B. Vipond⁵, M.C. Eisler¹

¹Bristol Veterinary School, University of Bristol, Langford, ²Rothamsted Research, North Wyke, ³Rothamsted Research, Harpenden, ⁴University of Bristol, Bristol Children's Vaccine Centre, Bristol, ⁵Public Health Laboratory Bristol, Public Health England, Bristol

amyc.thomas@bristol.ac.uk

Background

In both humans and livestock, viral-bacterial interactions in the upper respiratory tract (URT) appear to be important for development of respiratory disease. In healthy pre-school children, density of commensal nasopharyngeal bacteria has been demonstrated to increase following live-attenuated influenza vaccine. Dysbiosis of respiratory microbiota may influence transmission dynamics.

Objectives

To investigate carriage of *Pasteurellaceae* in the bovine URT and whether carriage density increases following mild respiratory viral infection.

Methods

Two cohorts of healthy beef suckler calves were bred, and housed post-weaning in three identical barns at the BBSRC National Capability, North Wyke Farm Platform (BBSRC, BB/J004308) over two successive years. We evaluated baseline carriage and density of *Pasteurellaceae* in calves born in 2015 (n = 60) and the effects of respiratory viral infection on *Pasteurellaceae* colonisation in calves born in 2016 (n = 87); using a stepped wedge design double-blind cluster randomised trial. Calf clusters (n = 29) received intranasal live-attenuated vaccine (RS+PI3) or placebo. Nasal swabs and jugular blood were collected from calves. Three qPCR assays were developed to detect and quantify *Pasteurella multocida*, *Mannheimia haemolytica* and *Histophilus somni*.

Results

The stepped wedge cluster-randomised trial will be described. Baseline rates of carriage were higher for *P. multocida* (57/60; CI_{95%}: 0.852 – 0.987 and 72/87; CI_{95%}: 0.728 – 0.897) than for *H. somni* (30/60; CI_{95%}: 0.370 – 0.630 and 36/87; CI_{95%}: 0.311 – 0.524) in both years. In absence of vaccination, carriage for both species decreased over time. Density of *P. multocida* carriage at baseline ranged widely over 5 orders of magnitude.

Conclusion

We have developed methods for rapid detection and quantification of *Pasteurellaceae* from nasal swabs, and demonstrated their usefulness in investigating pathogen colonisation of the bovine URT. We are currently using these methods together with attenuated vaccine viruses as tools to explore respiratory host-microbiome interactions in cattle.